

(k, m) -Catalan Numbers and Hook Length Polynomials for Plane Trees

Rosena R.X. Du^{*} and Fu Liu[†]

February 1, 2008

Abstract: Motivated by a formula of A. Postnikov relating binary trees, we define the hook length polynomials for m -ary trees and plane forests, and show that these polynomials have a simple binomial expression. An integer value of this expression is $C_{k,m}(n) = \frac{1}{mn+1} \binom{(mn+1)k}{n}$, which we call the (k, m) -Catalan number. For proving the hook length formulas, we also introduce a combinatorial family, (k, m) -ary trees, which are counted by the (k, m) -Catalan numbers.

Keywords: (k, m) -Catalan number, (k, m) -ary tree, binary tree, m -ary tree, plane forest, hook length polynomial.

AMS Classification: 05A15, 05A19.

1 Introduction

The main result in this paper is originally motivated by seeking a simple bijective proof of the following identity:

$$\sum_T \frac{n!}{2^n} \prod_v \left(1 + \frac{1}{h_v}\right) = (n+1)^{n-1}, \quad (1.1)$$

where the sum is over all complete binary trees with n internal vertices, the product is over all internal vertices of T , and h_v is the “hook length” of v in T , namely, the number of internal vertices in the subtree of T rooted at v . As an example, Figure 1 shows the five complete binary trees of three internal vertices, with each internal vertex labelled by its hook length. In this case, this identity says that $3 + 3 + 4 + 3 + 3 = (3 + 1)^2$.

This identity was first derived by Postnikov [6], who also asked for a combinatorial proof of this identity. Chen and Yang [2] and Seo [7] both gave bijective proofs of it.

Based on (1.1), Lascoux conjectured that if we substitute x for 1 on the left hand side, we

^{*}Department of Mathematics, East China Normal University, Shanghai, P. R. China, rxdu@math.ecnu.edu.cn.

[†]Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA, fuliu@math.mit.edu.




Figure 1: Hook lengths for the five binary trees with 3 internal vertices.

will get the following polynomial:

$$\sum_T \prod_v \left(x + \frac{1}{h_v} \right) = \frac{1}{(n+1)!} \prod_{i=0}^{n-1} ((n+1+i)x + n+1-i). \quad (1.2)$$

This is equivalent to the following, more suggestive identity:

$$\sum_T \prod_v \frac{(h_v + 1)x + 1 - h_v}{2h_v} = \frac{1}{n+1} \binom{(n+1)x}{n}. \quad (1.3)$$

We call the left side of (1.3) the “hook length polynomial” of complete binary trees. Note that if we replace x with k , the right hand side of (1.3) becomes $\frac{1}{n+1} \binom{(n+1)k}{n}$, which is exactly the number of complete k -ary trees with $n+1$ internal vertices. In fact we could prove (1.3) by showing that both the hook length polynomials for complete binary trees and k -ary trees have the same recurrence relation.

Moreover, we are able to generalize (1.3) from enumerating on binary trees to enumerating on $(m+1)$ -ary trees, where $m \geq 0$, and obtain the following identity for hook length polynomials for $(m+1)$ -ary trees:

$$\sum_T \prod_v \frac{(mh_v + 1)x + 1 - h_v}{(m+1)h_v} = \frac{1}{mn+1} \binom{(mn+1)x}{n}. \quad (1.4)$$

Identity (1.4) is one of the main results in this paper. If we replace x with k , the right hand side of (1.4) becomes

$$\frac{1}{mn+1} \binom{(mn+1)k}{n}.$$

We define this number as the (k, m) -Catalan number of order n . We want to prove (1.4) using the similar idea as for (1.3). Therefore, we need to find some nice combinatorial interpretation of this number.

In Section 2, we define a combinatorial structure, (k, m) -ary trees, and prove that they are counted by (k, m) -Catalan numbers. In Section 3, we prove (1.4) by showing that the hook length polynomials for $(m+1)$ -ary trees and (k, m) -ary trees have the same recurrence relation. Some other nice identities concerning hook lengths for $(m+1)$ -ary trees are also studied in this section. In Section 4, we give similar results for hook length polynomials for plane forests.

2 (k, m) -ary Trees and (k, m) -Catalan numbers

In this section we define (k, m) -ary trees, and prove that they are counted by (k, m) -Catalan numbers.

Let us first review some terminology related to trees. A *tree* is an acyclic connected graph, and a *forest* is a graph such that every connected component is a tree. In this paper we will assume all the trees are *unlabelled plane trees*, i.e., rooted trees whose vertices are considered to be indistinguishable, but the subtrees at any vertex are linearly ordered. For each vertex of a tree, we say that it is of *degree m* if it has m children, and call vertices of degree 0 *leaves*. (Note that the definition of degree here is different from that in graph-theoretic terminology.) Vertices that are not leaves are called *internal vertices*. For any tree T , we use $\mathcal{I}(T)$ to denote the set of internal vertices of T .

A *complete binary tree* is a tree each of whose internal vertices has degree 2, and a *complete m -ary tree* is a tree each of whose internal vertices has degree m . Since all of our trees will be complete, we will frequently just say m -ary tree or binary tree. We will use $\mathcal{T}_m(n)$ to denote the set of all m -ary trees with n internal vertices. An m -ary tree with 0 internal vertices is defined to be a single vertex.

Let T be a tree with root r . For each vertex v of T , we say that v is on *level j* if the unique path from v to r is of length j , and the root is said to be on the *level 0*. We can then define (k, m) -ary trees as follows.

Definition 2.1 *For $k, m \geq 1$ and $n \geq 0$, a (k, m) -ary tree of order n is a tree which satisfies the following:*

1. *All vertices on even levels have degree k .*
2. *All vertices on odd levels have degree m or 0, and there are exactly n vertices of degree m .*

For example, the tree in Figure 2 is a $(3, 2)$ -ary tree of order 3. We will use $\mathcal{T}_{k,m}(n)$ to denote the set of (k, m) -ary trees of order n . It is easy to check that there are a total of $(mn + 1)(k + 1)$ vertices for each tree $T \in \mathcal{T}_{k,m}(n)$, with $mn + 1$ vertices on even levels (which are all internal vertices), and $(mn + 1)k$ vertices on odd levels (in which there are n internal vertices and $(mn + 1)k - n$ leaves). We will call the n internal vertices on odd levels *crucial vertices*. Note that a (k, m) -ary tree of order 0 (with 0 crucial vertices) is a k -ary tree with only one internal vertex (which is the root).

Theorem 2.2 *The number of (k, m) -ary trees of order n is*

$$C_{k,m}(n) := \frac{1}{mn + 1} \binom{(mn + 1)k}{n}. \quad (2.1)$$

Proof. Let $C(z) = \sum_{i=0}^{\infty} c_i z^i$ be the generating function for (k, m) -ary trees, where c_n is the number of (k, m) -ary trees of order n .




Figure 2: A $(3, 2)$ -ary trees of order 3.

For $T \in \mathcal{T}_{k,m}(n)$, if we delete the root of T , we will get k subtrees, and each of them can be considered as a forest of m ordered (k, m) -ary trees. Hence we have

$$C = (1 + zC^m)^k. \quad (2.2)$$

Let $B = zC^m$, then from (2.2) we have $C = (1 + B)^k$ and $B = z(1 + B)^{mk}$. Applying the Lagrange inversion formula (see [9, p.42]) yields

$$[z^n]C(z) = \frac{1}{n}[t^{n-1}] \left(\frac{d}{dt}(1+t)^k \right) (1+t)^{mkn} = \frac{1}{mn+1} \binom{(mn+1)k}{n}.$$

■

Remarks:

1. We can also prove this theorem (as well as Lemma 4.10 in Section 4) combinatorially, but we omit these proofs for the sake of brevity. The current proofs of Theorem 2.2 and Lemma 4.10 were suggested by a referee.
2. It is well known [9, Ex. 6.19] that binary trees with n internal vertices are counted by the n th Catalan number:

$$|\mathcal{T}_2(n)| = C(n) = \frac{(2n)!}{(n+1)!n!} = \frac{1}{2n+1} \binom{2n+1}{n} = \frac{1}{n} \binom{2n}{n-1} = \frac{1}{n+1} \binom{2n}{n}, \quad (2.3)$$

and it is also known [3, 4, 5] that the number of m -ary trees with n internal vertices is

$$|\mathcal{T}_m(n)| = C_m(n) = \frac{1}{n} \binom{mn}{n-1} = \frac{1}{mn+1} \binom{mn+1}{n}. \quad (2.4)$$

For a (k, m) -ary tree of order n , if k or m equals 1, we can simply contract the redundant edges and get the following results:

$$\begin{aligned} \mathcal{T}_{1,2}(n) &= \mathcal{T}_2(n), & \mathcal{T}_{2,1}(n) &= \mathcal{T}_2(n+1), \\ \mathcal{T}_{1,m}(n) &= \mathcal{T}_m(n), & \mathcal{T}_{k,1}(n) &= \mathcal{T}_k(n+1), \end{aligned}$$

which coincides with the fact that

$$C_{1,2}(n) = C(n), \quad C_{2,1}(n) = C(n+1), \quad (2.5)$$

$$C_{1,m}(n) = C_m(n), \quad C_{k,1}(n) = C_k(n+1). \quad (2.6)$$

Hence (k, m) -Catalan numbers $C_{k,m}(n)$ can be viewed as a generalization of $C(n)$ and $C_k(n)$, justifying the choice of terminology.

3 Hook length polynomials for m -ary trees

In this section we will study hook length polynomials for m -ary trees. Given an m -ary tree T , for any internal vertex v , recall that the *hook length* h_v of v is defined as the number of internal vertices in the subtree rooted at v .

We define the *hook length polynomial* for an m -ary tree T to be

$$T(x) = \prod_{v \in \mathcal{I}(T)} \frac{((m-1)h_v + 1)x + 1 - h_v}{mh_v}.$$

For $n \geq 1$, the n th *hook length polynomial of m -ary trees* is then defined to be

$$H_{n,m}(x) = \sum_{T \in \mathcal{T}_m(n)} T(x),$$

and by convention, we set $H_{0,m}(x) = 1$. (Recall that $\mathcal{T}_m(n)$ denotes the set of all m -ary trees with n internal vertices.)

The main result of this section is the following identity for the hook length polynomial of $(m+1)$ -ary trees:

Theorem 3.3

$$H_{n,m+1}(x) = \frac{1}{mn+1} \binom{(mn+1)x}{n}. \quad (3.1)$$

For the example in Figure 1, we have

$$\begin{aligned} T_1(x) &= T_2(x) = T_4(x) = T_5(x) = \frac{1}{12}x(2x-1)(3x-1), \\ T_3(x) &= \frac{1}{3}x^2(2x-1), \end{aligned}$$

and

$$H_{3,2}(x) = 4 \cdot \frac{1}{12}x(2x-1)(3x-1) + \frac{1}{3}x^2(2x-1) = \frac{1}{4} \binom{4x}{3},$$

which coincides with equation (3.1).

We will prove Theorem 3.3 by showing that both sides of (3.1) satisfy the same recurrence, as proved in the following two lemmas.

Lemma 3.4

$$H_{n,m+1}(x) = \frac{(mn+1)x + 1 - n}{(m+1)n} \sum_{\substack{i_1, i_2, \dots, i_{m+1} \geq 0 \\ i_1 + i_2 + \dots + i_{m+1} = n-1}} H_{i_1, m+1}(x) H_{i_2, m+1}(x) \cdots H_{i_{m+1}, m+1}(x). \quad (3.2)$$

Proof. For the sake of convenience, we set

$$h_v(x) = \frac{(mh_v + 1)x + 1 - h_v}{(m+1)h_v}.$$

For any $T \in \mathcal{T}_{m+1}(n)$, if r is the root of T , then $h_r = n$, so $h_r(x) = \frac{(mn+1)x+1-n}{(m+1)n}$. Let $T_1, T_2, \dots, T_{m+1}$ be the $(m+1)$ -ary trees obtained by deleting r from T , so the total number of internal vertices in $T_1, T_2, \dots, T_{m+1}$ is $n-1$, and

$$T(x) = \frac{(mn+1)x+1-n}{(m+1)n} T_1(x) T_2(x) \cdots T_{m+1}(x).$$

If T_j has i_j internal vertices for $1 \leq j \leq m+1$, by summing over all $T \in \mathcal{T}_{m+1}(n)$ we get the desired result. $\blacksquare$

Lemma 3.5

$$C_{k,m}(n) = \frac{(mn+1)k+1-n}{(m+1)n} \sum_{\substack{i_1, i_2, \dots, i_{m+1} \geq 0 \\ i_1 + i_2 + \dots + i_{m+1} = n-1}} C_{k,m}(i_1) C_{k,m}(i_2) \cdots C_{k,m}(i_{m+1}) \quad (3.3)$$

Proof. We will prove this recursion by considering the structure of (k, m) -ary trees. Let T be a tree in $\mathcal{T}_{k,m}(n)$ with one crucial vertex v circled; clearly there are $n C_{k,m}(n)$ such trees. If we delete the m edges immediately below v , we get a forest of m ordered (k, m) -ary trees $T_1, T_2, \dots, T_m$ together with a (k, m) -ary tree T_{m+1} which has a leaf circled. Suppose T_j has i_j crucial vertices for all $1 \leq j \leq m+1$, so $i_1 + i_2 + \dots + i_{m+1} = n-1$. It is easy to see that such a split operation is a bijection between trees in $\mathcal{T}_{k,m}(n)$ with one circled crucial vertex and the set of $(m+1)$ -tuples $(T_1, T_2, \dots, T_m; T_{m+1})$ such that $T_1, T_2, \dots, T_m$ are linearly ordered and T_{m+1} has a circled leaf (see Figure 3), therefore we have

$$\#\{(T_1, T_2, \dots, T_m; T_{m+1})\} = n C_{k,m}(n). \quad (3.4)$$

Next we will finish the proof by showing that the number of such tuples equals n times the right hand side of (3.3). Let $(T'_1, T'_2, \dots, T'_m, T'_{m+1})$ be any set of $m+1$ ordered (k, m) -ary trees with a total of $n-1$ crucial vertices such that there is one tree in the set (not required to be T'_{m+1}) which has a circled leaf. Since a (k, m) -ary tree with i_j crucial vertices has $(mi_j+1)k-i_j$ leaves, there are $(mn+1)k+1-n$ leaves among all these trees. Hence we have

$$\#\{(T'_1, T'_2, \dots, T'_m, T'_{m+1})\} = ((mn+1)k+1-n) \cdot \sum_{\substack{i_1, i_2, \dots, i_{m+1} \geq 0 \\ i_1 + i_2 + \dots + i_{m+1} = n-1}} C_{k,m}(i_1) C_{k,m}(i_2) \cdots C_{k,m}(i_{m+1}). \quad (3.5)$$

On the other hand, for each set $(T'_1, T'_2, \dots, T'_m, T'_{m+1})$, we can always set the tree which has a circled leaf to be the last one to form an $(m+1)$ -tuple $(T_1, T_2, \dots, T_m; T_{m+1})$, so

$$\#\{(T'_1, T'_2, \dots, T'_m, T'_{m+1})\} = (m+1) \#\{(T_1, T_2, \dots, T_m; T_{m+1})\}. \quad (3.6)$$

Compare (3.4), (3.5) and (3.6), we get (3.3). $\blacksquare$

Proof of Theorem 3.3: Since $H_{n,m+1}(x)$ and $\frac{1}{mn+1} \binom{(mn+1)x}{n}$ are both polynomials of x of degree n , it is enough to prove that both sides coincide for positive integer values of x . For any positive integer k , we have

$$C_{k,m}(0) = \binom{k}{0} = 1 = H_{0,m+1}(k).$$




Figure 3: A $(3,2)$ -tree of order 3 with one circled crucial vertex and the corresponding forest $(T_1, T_2; T_3)$, where (T_1, T_2) is an ordered pair and T_3 has a circled leaf.

Moreover, according to Lemma 3.4 and Lemma 3.5, $H_{n,m+1}(k)$ and $C_{k,m}(n)$ have the same recurrence relation. Therefore for any $n \geq 0$, we have

$$H_{n,m+1}(k) = C_{k,m}(n), \forall k \in \mathbb{N}.$$

Hence we have finished the proof. ■

Corollary 3.6

$$\sum_{T \in \mathcal{T}_{m+1}(n)} \prod_{v \in \mathcal{I}(T)} \left(x + \frac{1}{h_v} \right) = \frac{1}{(mn+1)n!} \prod_{i=0}^{n-1} ((mn+1+i)x + mn+1-mi). \quad (3.7)$$

Proof. Define φ_n, E and R_m to be the functions that map polynomials of degree at most n to polynomials of degree at most n , given by $\varphi_n(f(x)) = x^n f(1/x)$, $E(f(x)) = f(x+1)$, $E^{-1}(f(x)) = f(x-1)$, $R_m(f(x)) = f((m+1)x+m)$, for any polynomial $f(x)$. Identity (3.7) can then be obtained by applying $E^{-(m-1)} \circ \varphi_n \circ E^{-1} \circ \varphi_n \circ R_m$ to both sides of (3.1). ■

If we choose the special values 0 or m for x in (3.7), we will get the following identities.

Corollary 3.7

$$n! \sum_{T \in \mathcal{T}_{m+1}(n)} \prod_{v \in \mathcal{I}(T)} \frac{1}{h_v} = \prod_{i=0}^{n-1} (mi+1), \quad (3.8)$$

$$\sum_{T \in \mathcal{T}_{m+1}(n)} \prod_{v \in \mathcal{I}(T)} \left(m + \frac{1}{h_v} \right) = \frac{(m+1)^n (mn+1)^{n-1}}{n!}. \quad (3.9)$$

We want to remark here that equation (3.8) gives the number of increasing $(m+1)$ -ary trees and is also proved by Stanley in [8].

4 Hook length polynomials for plane forests

We define a *plane forest* to be a forest of plane trees which are linearly ordered, and use $\mathcal{F}(n)$ to denote the set of plane forests with n vertices. Given a plane forest $F \in \mathcal{F}(n)$, for any vertex v of F , the *hook length* h_v of v is defined as the number of vertices in the subtree rooted at v . (Note that this definition is slightly different from the definition of hook length for m -ary trees. For the latter one, we count the number of *internal* vertices because we make our m -ary trees complete, and the definition in terms of internal vertices is more natural in this context. But we prefer to considering complete m -ary trees in this paper in order to make the definition of (k, m) -ary trees and many proofs clearer.)

In this section we will study the hook length polynomial for plane forests.

The *hook length polynomial* for a plane forest F is defined to be

$$F(x) = \prod_{v \in V(F)} \frac{(2h_v - 1)x + 1 - h_v}{h_v},$$

where $V(F)$ is the set of vertices of F .

The n th *hook length polynomial of plane forests* is then defined to be

$$H_n(x) = \sum_{F \in \mathcal{F}(n)} F(x).$$

By convention, we set $H_0(x) = 1$.

The main result of this section is the following identity for the hook length polynomial of plane forests:

Theorem 4.8

$$H_n(x) = \frac{1}{2n+1} \binom{(2n+1)x}{n}. \quad (4.1)$$

As in the last section, we will prove (4.1) by showing that both sides satisfy the same recurrence for integer values of x .

Lemma 4.9

$$H_n(x) = \sum_{i=1}^n \frac{(2i-1)x - (i-1)}{i} H_{i-1}(x) H_{n-i}(x). \quad (4.2)$$

Proof. For $n \geq 1$, given any plane forest $F \in \mathcal{F}(n)$, we pick the first plane tree T of F . Suppose T has i vertices, so the rest of the trees in F can be considered as a plane forest with $n-i$ vertices. Let $V(T)$ be the set of vertices of T and r the root of T , so $h_r = i$. By removing

r from T , we get another plane forest which has $i - 1$ trees. Therefore,

$$\begin{aligned}
H_n(x) &= \sum_{i=1}^n \left(\prod_{v \in V(T)} \frac{(2h_v - 1)x + 1 - h_v}{h_v} \right) H_{n-i}(x) \\
&= \sum_{i=1}^n \frac{(2h_r - 1)x + 1 - h_r}{h_r} \left(\prod_{v \in V(T), v \neq r} \frac{(2h_v - 1)x + 1 - h_v}{h_v} \right) H_{n-i}(x) \\
&= \sum_{i=1}^n \frac{(2i - 1)x + 1 - i}{i} \left(\prod_{v \in V(T), v \neq r} \frac{(2h_v - 1)x + 1 - h_v}{h_v} \right) H_{n-i}(x) \\
&= \sum_{i=1}^n \frac{(2i - 1)x + 1 - i}{i} H_{i-1}(x) H_{n-i}(x).
\end{aligned}$$

■

For the right side of (4.1), if we replace x with k , we will get $\frac{1}{2n+1} \binom{(2n+1)k}{n} = C_{k,2}(n)$. When $k = 1$, as we mentioned before $C_{1,2}(n) = C(n)$ is just a Catalan number, which satisfies the recurrence relation:

$$C(n) = \sum_{i=1}^n C(i-1)C(n-i).$$

The following lemma gives a generalization of this recurrence relation.

Lemma 4.10 *For $n \geq 1$,*

$$C_{k,2}(n) = \sum_{i=1}^n \frac{(2i-1)k - (i-1)}{i} C_{k,2}(i-1) C_{k,2}(n-i). \quad (4.3)$$

Proof. As in the proof of Theorem 2.2, we let $C(z)$ be the generating function for $(k, 2)$ -ary trees. And we let $D(z) = z \sum_{i=0}^{\infty} \frac{c_i}{i+1} z^i$ be $C(z)$'s antiderivative. By (2.2),

$$C = (1 + zC^2)^k.$$

Differentiating on both sides, we get

$$\begin{aligned}
C' &= k(1 + zC^2)^{k-1} (C^2 + 2zCC') \\
&= k \frac{C}{1 + zC^2} (C^2 + 2zCC').
\end{aligned}$$

One can rearrange this equation to obtain

$$(1 - k)C = -(2k - 1)(C + zC') + \frac{C'}{C^2}.$$

Then integrating on both sides gives us

$$(1 - k)D = -(2k - 1)zC - \frac{1}{C} + \alpha,$$

where α is a constant. If we let $z = 0$, we find that $\alpha = 1$. The former equation can be written as

$$\frac{C-1}{z} = (2k-1)C^2 + (1-k)C\frac{D}{z}. \quad (4.4)$$

We complete the proof by observing that (4.3) and (4.4) are equivalent. $\blacksquare$

Proof of Theorem 4.8: For any positive integer k , we have:

$$H_0(k) = 1 = \frac{1}{2 \cdot 0 + 1} \binom{(2 \cdot 0 + 1)k}{0} = C_{k,2}(0).$$

According to Lemma 4.9 and Lemma 4.10, $C_{k,2}(n)$ and $H_n(k)$ satisfy the same recurrence relation. Therefore, for any nonnegative integer n ,

$$\frac{1}{2n+1} \binom{(2n+1)k}{n} = C_{k,2}(n) = H_n(k), \forall k \in \mathbb{N}.$$

Since both $\frac{1}{2n+1} \binom{(2n+1)x}{n}$ and $H_n(x)$ are polynomials of degree n and they agree on infinitely many values of x , we have proved the desired result. $\blacksquare$

Similar to the proof of Corollary 3.6, if we apply $E \circ \varphi_n \circ E \circ \varphi_n$ to both sides of (4.1), we get the following identity.

Corollary 4.11

$$\sum_{F \in \mathcal{F}(n)} \prod_{v \in V(F)} \left(x + \frac{1}{h_v} \right) = \frac{1}{(2n+1)n!} \prod_{i=0}^{n-1} ((2n+1-i)x + (2n+1-2i)). \quad (4.5)$$

Finally, setting $x = 0$ or -2 , we obtain two additional interesting formulas.

Corollary 4.12

$$n! \sum_{F \in \mathcal{F}(n)} \prod_{v \in V(F)} \frac{1}{h_v} = (2n-1)!!, \quad (4.6)$$

$$\sum_{F \in \mathcal{F}(n)} \prod_{v \in V(F)} \left(2 - \frac{1}{h_v} \right) = \frac{(2n+1)^{n-1}}{n!}. \quad (4.7)$$

Remark: By comparing (3.1) and (4.1), one may have noticed that the hook length polynomials for ternary trees and plane forests are the same:

$$H_{n,3}(x) = F_n(x). \quad (4.8)$$

In fact, when $m = 2$, the right hand side of (3.7) and (4.5) differ only in one sign and agree with each other when $x = 0$. This reflects the fact that increasing ternary trees with n internal vertices and increasing plane forests on n vertices are both counted by $(2n-1)!!$, as proved in [8] and [1], respectively. We wonder if there is any direct explanation for (4.8).

Acknowledgments. The authors would like to thank Prof. Alain Lascoux for his encouragement in working on this topic, and who conjectured the special case $m = 1$ of Corollary 3.6.

They would also like to thank Prof. Richard P. Stanley for helpful discussions and suggestions, and two referees for their very valuable comments. Most of this work was done during the first author's visit to the Department of Mathematics of Massachusetts Institute of Technology, and she would also like to thank Prof. Stanley for his advice and support.

References

- [1] W. Y. C. Chen, A general bijective algorithm for increasing trees, *Systems Sci. Math. Sci.* 12 (1999), no. 3, 193–203.
- [2] W. Y. C. Chen, L. L. M. Yang, On the hook length formula for binary trees, preprint.
- [3] I. P. Goulden and D. M. Jackson, *Combinatorial Enumeration*, John Wiley, New York, 1983.
- [4] D. A. Klarner, Correspondences between plane trees and binary sequences, *J. Combinatorial Theory, Series A*. 9 (1970), 401–411.
- [5] D. E. Knuth, *The Art of Computer Programming*, Vol. 1, Fundamental Algorithms, Addison-Wesley Publishing Company, 1968.
- [6] A. Postnikov, Permutohedra, associahedra, and beyond. Stanley Conference, June 26, 2004, MIT. Slides available at: <http://www-math.mit.edu/~apost/talks/perm-slides.pdf>
- [7] S. Seo, A combinatorial proof of Postnikov's identity and a generalized enumeration of labeled trees, arXiv: math.CO/0409323 v1.
- [8] R. P. Stanley, Ordered Structures and Partitions, *Memoirs of the Amer. Math. Soc.*, no. 119 (1972).
- [9] R. P. Stanley, *Enumerative Combinatorics*, Vol. 2, Cambridge University Press, Cambridge, UK, 1999.